

2021 UPDATE



TEXAS A&M UNIVERSITY

Office of the Texas State
Climatologist

ASSESSMENT of HISTORIC
and FUTURE TRENDS of
EXTREME WEATHER
IN TEXAS, 1900-2036



Cover photo by John Nielsen-Gammon

AUTHOR TEAM:

John Nielsen-Gammon*, Sara Holman*, Austin Buley*, Savannah Jorgensen*

ADDITIONAL AUTHORS, ORIGINAL REPORT:

Jacob Escobedo*, Catherine Ott[^], Jeremy Dedrick*, Ali Van Fleet*

*Department of Atmospheric Sciences, Texas A&M University

[^]Environmental Programs, College of Geosciences, Texas A&M University

OCTOBER 7, 2021, DOCUMENT OSC-202101

EXECUTIVE SUMMARY

This report analyzes historic observations of temperature, precipitation, and extreme weather in Texas and identifies ongoing and likely future trends out to the year 2036. These trends represent climatological expectations; the actual weather from year to year and decade to decade will be heavily influenced by natural variability which at this point is largely unpredictable.

The average annual Texas surface temperature in 2036 is expected to be 3.0 °F warmer than the 1950-1999 average and 1.8 °F warmer than the 1991-2020 average. The number of 100-degree days at typical stations is expected to nearly double by 2036 compared to 2001-2020, with higher frequency of 100-degree days in urban areas. Extreme monthly summertime temperature trends imply an increase of about 1 °F compared to the 1950-1999 average. Meanwhile, extreme monthly wintertime temperatures are expected to continue to increase at an even faster rate, and the coolest days of the summer are expected to continue becoming warmer.

Texas precipitation has increased by 10% or more in eastern Texas but little trend is present in western Texas. Precipitation trends to 2036 are likely to be dominated by natural variability. Extreme precipitation is expected to increase in intensity on average statewide by 6%-10% relative to 1950-1999 and 2%-3% relative to 2001-2020. This translates to an increase in the frequency of extreme rain of 30%-50% relative to the climatological expected frequency in 1950-1999 and 10%-15% relative to 2001-2020. Historical trends so far represent an increase of extreme precipitation intensity by about 7% from 1960 to 2020.

Drought will continue to be driven largely by multidecadal precipitation variability, with long-term precipitation trends expected to be relatively small. Other factors affecting drought impacts, such as increased temperatures and improved plant water use efficiency, on balance decrease water availability but will cause drought impact trends to be highly sector-specific, with the impacts possibly smaller for agriculture than for surface water supply.

River flooding is subject to a similar mix of factors as drought. No long-term river flooding trend has been identified in the observations, nor is such a trend projected at this point, except perhaps for the most extreme floods and areas with normally high rainfall. Urban flooding is projected to increase, both as a simple matter of urban population increase and because of the projected increase of precipitation intensity, which drives flooding in fast-response drainages like those usually found in urban areas. The climate-driven trend in urban flood frequency should be similar to the climate-driven trend in extreme precipitation frequency: 30%-50% in 2036

relative to 1950-1999 and 10%-15% relative to 2001-2020.

Winter weather can be dangerous in Texas in part because it is relatively rare in most areas of the state. The frequency of extreme winter weather ought to decrease in Texas because the existence of winter weather is dependent on temperatures being cold enough to support winter weather. As the climate warms, the likelihood of winter weather decreases. Both extreme cold and snowfall either become less frequent or are expected to do so. Widespread snowfall events in Texas such as the one that took place in February 2021 are extremely rare.

Projections of severe weather associated with thunderstorms are difficult because the historical records are uneven and future thunderstorms are too small to be simulated with global climate models. Indirect evidence supports an increase in the number of days capable of producing severe thunderstorms and an increase in the frequency of very large hail in early springtime, but these possible trends are too uncertain to quantify.

The combination of coastal subsidence and sea level rise is contributing to or driving a general retreat of the Texas coastline, both along the barrier islands and in coastal wetlands. Relative sea level rise is expected to continue at similar average rates in the near future, as reduced groundwater extraction is balanced by accelerating sea level rise. Storm surges from hurricanes will tend to be more severe because of higher relative sea levels, and a possible increase in extreme hurricane intensity may further increase storm surge risk.

Weather and climate drivers of wildfire risk are projected to increase the risk of wildfires throughout the state, primarily due to increased rates of drying and increased fuel load. The increase in wildfire risk may not be as large in far West Texas where rising temperatures and decreasing precipitation may overcome the carbon dioxide fertilization effect and lead to less accumulation of fuels.

Funding for this report was provided by Texas 2036. We thank Andrew Dessler and three anonymous reviews, which helped to improve the content of the original report, OSC-202001.

Suggested citation:
Nielsen-Gammon, J., S. Holman, A. Buley, S. Jorgensen, J. Escobedo, C. Ott, J. Dedrick, and A. Van Fleet, 2021: *Assessment of Historic and Future Trends of Extreme Weather in Texas, 1900-2036: 2021 Update*. Document OSC-202101, Office of the State Climatologist, Texas A&M University, College Station, 44 pp.

INTRODUCTION



Texas is vulnerable to a wide range of natural hazards, most of which are associated with weather and climate events. The natural environment has evolved partly in response to these natural hazards. For example, plant hardiness is largely determined by ability to survive extreme winter cold and drought. The built environment, including for example homes, roads, and power plants, is designed to a certain level of resiliency to natural hazards. Human activities as fundamental to survival as food production and water supply are tailored to the particular combination of weather and climate risks at play in a given location.

The future of Texas depends on its resilience to the natural hazards of the future. It is up to Texans, both individually and collectively, to decide what level of resilience is appropriate, and at what cost, compared to the costs of damage and recovery on both an economic and societal level. Nobody knows which specific weather and climate events will befall Texas over the next couple of decades. But a wide variety of information can be used to estimate the risks of certain types of weather and climate events over that period.

The standard practice for estimating the risk of natural hazards has been to assume that future risk is equal to historical risk. This practice works if the underlying climatic conditions are unchanging. However, Texas climate is affected by changing patterns of vegetation, irrigation, and urbanization. Texas climate is also embedded in the global climate system, which is itself changing. All of these factors have influenced historical trends in weather and climate extremes and will continue to influence trends in the future. Given a changing climate, historic trends may provide a better guide to future risk than mere historical averages.

The sponsors of these projections requested that projections be based primarily on existing trends. Doing so makes sense only if the causes of those trends are understood and are expected to continue. The scientific understanding of the causes of trends draws upon a large body of research, utilizing both observations and experiments with global climate models. This report presents trends for a variety of historic periods, and the projections are based on historic trends that are expected to continue according to currently available science.¹

Several factors influence our ability to project historical trends into the future. First, historical data may not be sufficiently accurate or consistent over time to yield reliable trends. Second, natural climate variability and the randomness of extreme weather events can mask or even overwhelm any

underlying long-term trends. Despite these limitations, there are sound reasons to expect continued change in a variety of aspects of extreme Texas weather, and knowledge of such likely changes can be very useful in a variety of planning contexts.


Trends are not the only potentially useful historic information. Some natural climate variations that occur on multidecadal time scales have a substantial effect on Texas weather. The present-day scientific ability to accurately predict such variations twenty years into the future is quite limited. Nonetheless, knowledge of these variations informs the understanding of past trends and suggests whether recent weather patterns are representative of typical future conditions.

This report addresses historical and future trends in extreme temperatures, extreme precipitation, severe thunderstorms, and hurricanes. It also addresses trends in drought, floods, wildfire, and coastal erosion, to the extent that these natural hazards are affected by changes in weather and climate. For each natural hazard, the report considers the quality of the historical data, the historical risk and trends (data permitting), the causes of any observed or expected trends, and the projection of trends of future risk. For context, this report also considers trends in annual average temperature and precipitation.

Expected typical conditions in 2036 are expressed as a change compared to average conditions in 1950-1999 and 1991-2020. The latter period corresponds to the “normals” period as defined by the National Centers for Environmental Information (NCEI). When weather is reported as being x degrees or x inches above or below normal in weather forecasts or reports, it is relative to the 1991-2020 period.

With all projections, there is considerable uncertainty as to how things will actually turn out. This report does not attempt to quantify that uncertainty; prudent planning recognizes that we cannot know whether reality will end up higher or lower than the best available present-day estimates.

This report was commissioned and sponsored by Texas 2036. The report content is solely the responsibility of the authors. A previous version of this report has been peer-reviewed; reviewer comments and responses are available from the Office of the State Climatologist. This updated report is based on data and published research through September 2021. The analyses that are original to this report are based on data from publicly accessible data sources; analysis spreadsheets and software are available from the Office of the State Climatologist.



AVERAGE TEMPERATURES

While average temperatures do not themselves constitute weather and climate extremes, changes in average temperatures, either locally or globally, affect many aspects of extreme weather and climate trends. In addition, all else being equal, a change in average temperature would lead to a change in frequency of extreme temperatures, increasing hot extremes and decreasing cold extremes.

The National Centers for Environmental Information maintain very good analyses of monthly averages of daily maximum and minimum temperatures from 1895 to present throughout the lower 48 states.² The annual average daily maximum and minimum temperatures in Texas over the period 1897-2020 exhibit year-to-year variations of 2 °F or more. Broadly speaking, Texas temperatures climbed slightly during the early part of the 20th century, declined somewhat until the 1970s, and rose thereafter.

The rate of temperature increase since 1895 has averaged 0.11 °F per decade, less than the global average of 0.16 °F per decade.³ Indeed, the southeastern United States, including eastern Texas, is almost the only land area on the globe whose temperature increase over the 20th century was nearly zero.⁴ More recently, the Texas temperature trend has been larger. Since 1950, the trend is 0.27 °F per decade, and since 1975, 0.61 °F per decade. The global trend since 1975 is 0.36 °F per decade. Recent temperatures have increased in all seasons and in all regions of Texas.

The historic Texas temperature trend simulated by global climate models since 1950 is 0.32 °F per decade, and


since 1975, 0.55 °F per decade.⁵ About 90% of models simulate a 1975-2020 trend between 0.25 and 0.88 °F per decade. The simulated current rate of increase in Texas, based on the average of climate model projections for 2020-2040 for the low-emissions pathway RCP 4.5⁶, is around 0.62 °F per decade. Up to mid-century, climate projections are not very sensitive to the choice of emissions pathway.⁷

On the whole, the agreement between models and observations is decent. Factors that cause observed trends to differ from simulated trends include inadequacies in the models, inaccuracies in observations, actual and simulated natural variability, and local land surface changes such as irrigation and afforestation.


Historical data and climate models lead to similar conclusions. If recent trends continue, as expected, a middle-of-the-road estimate of the overall rate of temperature increase in Texas would be about 0.6 °F per decade. This means that **average Texas temperatures in 2036 should be expected to be about 1.8 °F warmer than the 1991-2020 average and 3.0 °F warmer than the 1950-1999 average. This would make a typical year around 2036 warmer than all but the absolute warmest year experienced in Texas during 1895-2020.**⁸ Even a very conservative extrapolation, based on the average of the 1950-2020 and 1975-2020 trends, would make a typical year around 2036 warmer than all but the five warmest years on record so far.

Temperature trends have been generally similar from season to season. During the period 1950-2020, summertime trends were smaller than trends in other seasons. During the period 1975-2020, wintertime trends were larger than trends in other seasons.


Average Temperature Trend, 1975-2020



Temperature trends since 1975 according to NCEI nClimDiv data.



Texas annual average statewide temperatures (°F)



Comparison of observed and simulated temperature trends in Texas



EXTREME TEMPERATURES

The projected changes in average temperature imply changes in unusually high or low temperatures as well. This assessment of extreme temperatures relies on two aggregated sets of temperature data: a set of stable stations (hereafter referred to as index stations)⁹ and a set with one composite station per county.¹⁰

To analyze actual trends in 100 °F days in Texas, the index stations were grouped into four regions. The analysis of changes in 100 °F days uses index stations grouped into four regions and analyzed collectively within each region.¹¹ Over the past 45 years, the linear trend shows an approximate doubling of the number of triple-digit days at stations in all four regions. Given past and projected temperature trends, a near-doubling of the number of 100-degree days between 2001-2020 and 2036 appears to be a reasonable projection.

Although the number of truly urban stations in Texas is limited, the existence of urban heat islands has likely led to an enhancement of 100 °F days in urban areas. Presently, the limited data suggests that triple-digit days are rising at similar rates in urban and rural areas.


An alternative way of examining extreme heat trends is to consider the average hottest day in each month during June-September in each county in Texas.¹² This metric is more sensitive to exceptional heat events than to sustained heat. It also enables us to use longer historical records, as this metric is not sensitive to changes in observing time.

Looked at this way, there is a very slight downward trend in extreme heat over the long term. This trend reflects the general pattern over the continental United States, where the most exceptional heat occurred back in the


1930s, with this summer across the United States being in a virtual tie with 1936 for warmest summer on record.¹³ The cause of this downward trend, which is in contrast to mean and extreme temperatures in all other seasons, may be related to the expansion of irrigation in the mid-20th century.¹⁴ Whatever the cause of the long-term slight downward trend, it appears to have reversed, with a substantial increase in extreme monthly heat in recent decades. Recent extreme monthly high temperatures have tended to be higher than the historic mean.

Extreme low temperatures during the winter months exhibit a stronger and more robust trend.¹⁵ Despite year-to-year fluctuations that are much larger for extreme cold than extreme heat, there is a long-term warming trend in monthly extreme cold temperatures across Texas, and the trend is larger in recent decades. Although they are much noisier, the absolute coldest temperature during winter shows similar long-term variations, with the extreme cold during 2021 easily identifiable.


No matter the metric, extreme cold is no longer as extreme as it used to be. This is broadly consistent with expectations: extreme cold air comes from the Arctic, which in general is warming faster than other parts of the globe. There have been some studies in recent years debating whether loss of Arctic sea ice and overall Arctic warming leads to changes in weather patterns that favor more intense incursions of Arctic air¹⁶, but this tendency, if present, has not been strong enough in Texas to stop the accelerating rise of extreme cold temperatures. This warming of extreme cold temperatures in Texas, much more rapid than warming of average temperatures, is consistent with what has been observed elsewhere in the Northern Hemisphere and is inconsistent with the argument that loss of Arctic sea ice is enhancing extreme cold over the continents.¹⁷



Trends and historic variability in index temperature stations in Texas, grouped by amount of urbanization. Arrows at top indicate data points that are above the margin of the graph. Trend lines are fit to the logarithm of 100 °F day counts to ensure non-negative values.




Hottest day of the month, June-September, at composite county stations across Texas, through 2021.



Average lowest temperature of the month, December-February, at composite county stations across Texas, and average lowest temperature overall.

Overall, extreme heat has recently become more frequent and more severe, while extreme cold has become less frequent and less severe overall. Trends in extreme cold are much larger than trends in extreme heat, which is leading to an overall decrease in the range of annual temperature extremes over time. That trend is also apparent within summer, as the lowest temperatures during July and August have become substantially warmer. Recent trends in the coolest temperatures of the summer are similar to recent trends in the hottest temperatures of the summer, so this reduction of temperature range may have ended.

Extreme summer heat is approaching values not seen since the early part of the 20th Century and is likely to surpass them by 2036. The typical number of triple-digit days by 2036 is projected to be substantially larger, about 40% larger than typical values so far in the 21st Century. **Meanwhile, the expected warming of extreme wintertime temperatures would make typical wintertime extremes by 2036 milder than most of the winter extremes in the historic record.** That said, extreme wintertime cold is more variable than other types of temperature extremes, so massively cold winter temperatures will continue to be possible.



Average lowest temperature of the month, July-August.

PRECIPITATION


Precipitation in Texas is quite variable, both in space and time. Much of the state has two rainy seasons, with the rainiest months on average being May, June, September, and October. In far West Texas, the wettest months are July and August, while far East Texas averages similar amounts in every month. Rainfall amounts increase from west to east, with the southeast corner of the state near Beaumont averaging over eight times the annual rainfall of some areas near El Paso.

The long-term trend of precipitation in Texas has been positive. Over the past century, parts of central and eastern Texas have experienced precipitation increases of 15% or more, while in much of the western part of the state the long-term trend is flat or even downward.¹⁸ The tendency for increasing precipitation in Texas


is not consistent with the majority of global climate models, with the average simulated trend being -2.6% per century.¹⁹ Models and observations both tend to feature more positive (or less negative) trends toward northeastern Texas than toward southwestern Texas.²⁰

Superimposed on the generally upward precipitation trend is considerable variability. El Niño has a prominent influence on cool-season rainfall in Texas: during El Niño years, Texas tends to be wet, while during La Niña years, Texas tends to be dry. Individual El Niño or La Niña events tend to last only 1-3 years, so this relationship is not useful for multidecadal predictions. However, both the Atlantic and Pacific Oceans feature multidecadal variability: the Atlantic Multidecadal Oscillation (AMO) and the Pacific Decadal Oscillation (PDO). Both have been shown to substantially influence precipitation in the United States, including Texas.²¹

Overall Precipitation Trend, 1895-2020



Precipitation trends since 1895 according to NCEI nClimDiv data.



Solid: Average Texas precipitation, inches, nine-year running averages, for March-June (green), July-October (orange), and November-February (gray), with dots showing values for last nine individual years. Dashed light green: Pacific Decadal Oscillation, November-June, nine-year running average, with squares showing values for last nine individual years. Dashed dark red: Atlantic Multidecadal Oscillation, March-October, sign reversed, nine-year running average, with squares showing values for last nine individual years. Both AMO and PDO have been rescaled and offset for ease of comparison with Texas precipitation.²²

The graph of decadal-scale variations of average Texas precipitation illustrates the distinct seasons during which the AMO and PDO influence Texas precipitation. The AMO was relatively high in the 1930s, 1950s, and 2000s and low in the 1900s, 1970s, 1980s, and early 1990s. The negative of the AMO is plotted here so that upticks in the graph correspond to upticks in Texas precipitation. The AMO pattern can be detected in the March-June precipitation (correlation = -0.25) and July-October precipitation (correlation = -0.47). Meanwhile, the PDO was relatively low near 1920, 1950, and 2010, and relatively high near 1940, 1980, and 1990. This pattern can be detected in the November-February precipitation (correlation = 0.53) and March-June precipitation (correlation = 0.54). So the AMO influences Texas precipitation on a decadal scale from spring to fall, while the PDO influences Texas precipitation from late fall through spring.

The decadal-scale variability is so large that the eye has difficulty detecting the aforementioned long-term upward trend in Texas precipitation. Over the next couple of

decades, this implies that variability is likely to have a far larger effect on precipitation changes than any long-term climate trend. Unfortunately, neither the AMO nor PDO can reliably be predicted two decades into the future. The last nine years illustrates that the PDO is much more erratic than the AMO.

Both the AMO and PDO favored unusually dry conditions in Texas during the first part of the 21st century. Since that pattern is unlikely to recur given the slow variability of the PDO and AMO, the expectation is that rainfall in Texas over the next two decades will tend to be greater than what was experienced from 1998 to 2012. The positive long-term precipitation trend further supports that expectation.

The climate model projections provide weak evidence for a precipitation decline. Unlike earlier model projections, the latest CMIP6 projections do not have precipitation in summertime declining more than precipitation in other seasons.²³



EXTREME RAINFALL


Many studies have documented an increase in extreme rainfall in Texas and surrounding areas for a variety of durations and thresholds.²⁴ On average across the region, extreme one-day precipitation has increased by 0.2" to 0.5" since the middle of the 20th century.²⁵ Within Texas, the local experience of extreme rainfall varies widely from place to place, with some locations having experienced a decrease in intensity of extreme rainfall over the period of data availability while the majority of locations have experienced an increase.²⁶ The median change since 1960 is an increase in intensity of about 7%.

All other things being equal, an increase in overall precipitation amounts would be expected to lead to an increase in extreme precipitation amounts, which inevitably implies an increase in the frequency of extreme precipitation above a given threshold. So the overall trend in Texas precipitation, discussed previously, contributes to the observed trend in extreme precipitation probabilities.

In addition to the overall precipitation effect, extreme rainfall is strongly affected by increased temperatures. A column of air that is producing rainfall will produce about 4% more rainfall for every °F of warming.²⁷ The extra precipitation intensity can also affect storm structure, and climate change can alter weather patterns and the frequency of dangerous storms. The direct temperature effect appears to be most important over the long haul in most midlatitude locations such as Texas. Changes in storm motion and frequency are expected to be comparatively subtle.²⁸

As noted earlier, computer model projections of overall rainfall amounts in Texas are somewhat inconsistent, but in general they show an overall leveling off or slight

1-day Duration 100-year Return Period Trends



Percentage change in the intensity of one-day rainfall with a 1% probability of occurring in any given year, based on a time-dependent statistical fit of annual daily precipitation maxima at composite stations.

decrease of precipitation amounts.²⁹ This suggests that trends of extreme precipitation in the future will be dominated by the increasing temperature effect and changes in storm structure. The strongest influence should be temperatures near the moisture source for Texas extreme precipitation, that is, the tropical oceans and the Gulf of Mexico. Tropical ocean temperatures are not expected to rise as quickly as temperatures over land in Texas, but would still be sufficient to produce an additional 2%-3% increase in extreme rainfall intensity from the temperature effect alone.³⁰ So extreme rainfall intensity and frequency are projected to continue increasing, though probably not as rapidly as they have increased in the past.

Rainfall risk is often characterized in terms of the 100-year rainfall event, which is an amount of rain over a given duration that has a 1% chance of occurring in any given year. If extreme rainfall amounts increase by just 20%, the 100-year rainfall event threshold is exceeded twice as often.³¹ So Texas's 7% median increase in the 100-year rainfall amount between 1960 and 2020 corresponds to a roughly 30% increase in the frequency of heavy rainfall exceeding the older 100-year threshold.

Based on projected temperatures and the dominance of the direct temperature effect on extreme rainfall, we anticipate an additional increase of 2%-3% in expected extreme rainfall intensity in 2036 compared to 2001-2020 and an overall increase of 6%-10% compared to 1950-1999. These changes in amount correspond to increases in the odds of extreme precipitation of 10%-15% and 30%-50%, respectively.

Note that the variations of extreme rainfall trends across Texas imply that one should not assume that recent extreme rainfall history in a given location is a suitable baseline for projecting future trends. While estimates of extreme rainfall risk based on historical data show a large uptick in Houston but less change in Dallas-Fort Worth³², climate change should be acting to increase the risk more uniformly across the entire state. The projected increase of extreme rainfall risk given above is relative to the expected past risk of extreme rainfall in a given location, not the actual occurrences of extreme rainfall in that location. Likewise, the actual extreme rainfall over the next few decades at any given location may defy the odds, either favorably or unfavorably.




DROUGHT


At first glance, the long-term increase in average rainfall should imply a decrease in drought as well. The linear trend in total statewide precipitation is about 1%-2% per decade, depending on the period considered.³³

The corresponding drought severity based on precipitation alone, as measured by the cumulative 9-month Standardized Precipitation Index, does indicate declines in severity. The change in cumulative dryness is -5.3% per decade since 1896, -12.8% per decade since 1950, and +11.1% per decade since 1975. As noted with respect to precipitation, the change since 1950 is large in part because of the extended drought that occurred early in the 1950s. The rise of cumulative dryness since 1975, however, has happened despite a slight increase of precipitation over the same period.


The explanation for this difference can be seen in the statewide precipitation graph. Note that the period of time from 1965 to 1985 featured very little precipitation variability, with statewide average precipitation ranging between 22" and 35". Since then, precipitation has become more variable, with 2011 precipitation below 14" and 2015 precipitation above 41". Greater precipitation variability leads to more intense droughts even if the overall precipitation amount doesn't change.



Statewide average precipitation, from NCEI's nClimDiv dataset.



Statewide annual cumulative drought severity, assessed using the SPI precipitation-only drought index.³⁴



Statewide annual cumulative drought severity, assessed using the Palmer Drought Severity Index.

The historic record of precipitation variability is, itself, quite variable, with no clear trend over the period of record despite the recent uptick in variability. Climate models generally simulate an increase in interannual precipitation variability in Texas since 1950 but little or no change in interannual precipitation variability going forward.³⁵ The lack of future variability is itself anomalous compared to the rest of the Northern Hemisphere land areas, which tend to show an increase in variability in excess of the increase in average precipitation itself.³⁶ So the recent trend of increased interannual variability is probably not going to continue apace through 2036, unless average precipitation itself continues to increase, and the precipitation variability experienced over 2000-2018 is probably representative of what should be expected through 2036.

Drought severity and impacts, however, involve much more than a lack of precipitation. The exact combination of factors determining the severity and impacts of a particular drought depend on the particular crop, water supply, or other system being affected, making it challenging to generalize trends in drought severity from conventional drought indices.³⁷

Temperature affects drought directly, by increasing the rate of evaporation from the soil and from water bodies. Many drought indices, such as the Palmer Drought Severity Index, attempt to include the effect of temperature on dryness.³⁸ Compared to rainfall-only SPI indices at various time scales, particularly the 9-month scale that the PDSI is most responsive to, the PDSI shows a less negative trend over the entire period and a larger positive trend recently.³⁹

However, while the SPI neglects the temperature effect on drought entirely, the conventional PDSI may overestimate it.⁴⁰ This is because, in addition to precipitation and temperature, carbon dioxide also affects drought. Elevated carbon dioxide levels improve the water use efficiency by plants, so would lead to increased soil moisture and decreased drought. Elevated carbon dioxide levels also increase biomass if plants are not otherwise water- or nutrient-limited, which might increase water use and decrease soil moisture. While these two effects work in opposite directions, the water use efficiency effect seems to be dominant, thereby allowing increased CO₂ to lead to improved plant growth despite meteorological factors leading to increased drought, at least on a global scale.⁴¹ Nonetheless, improved plant health overall does not eliminate possible detrimental effects from drought accompanied by increased temperatures, particularly since decreased plant water use itself implies increased temperatures.⁴² Also, different plants respond to carbon dioxide (and heat stress) differently.⁴³ Lastly, global climate model simulations that attempt to incorporate changes in plant physiology still indicate a substantial decrease in future soil moisture across the Great Plains.⁴⁴

While recent SPI values are near the long-term average, the SPI record indicates that much more severe precipitation deficits have occurred in the past. With warmer temperatures in recent decades, the recent series of droughts had impacts comparable to the droughts in the early and middle 20th century, despite smaller long-term precipitation deficits. This means that if (or when) another period of lack of precipitation develops like those in the early and middle 20th century, the higher



temperatures will lead to unprecedentedly severe drought impacts.

Increased carbon dioxide does not reduce the temperature effect of evaporation from lakes and reservoirs or from bare soil. Historically, there has been an increase in the evaporative capacity of the atmosphere across most of Texas, especially in West Texas and the Panhandle, and that trend is expected to continue due to robust projections of rising temperatures.⁴⁵ A continuation of the observed trend would lead to a roughly 4% increase in expected summertime evaporative losses from reservoirs in 2036 compared to 2000-2018.⁴⁶

For hydrological drought, runoff is important. Runoff changes depend on changes in soil moisture and changes in precipitation amount and intensity. Across the midlatitudes, there is on average a decrease of topsoil moisture but little change in deep soil moisture.⁴⁷ This would favor a slight decrease in runoff for a given amount of precipitation. Countering that is the tendency for greater rainfall intensity,⁴⁸ which would lead to a greater runoff fraction and in turn would partially cause the decrease in soil moisture, while a model-simulated change in precipitation seasonality in some areas could lead to greater drying in summer when drought is most severe.⁴⁹

Overall, the most detailed recent study of simulated streamflow impacts finds a projected increase in both drought severity and flood severity for the two studied

river basins. Those increases appear to be unaffected by errors in calculating the drought indices.⁵⁰

Despite all the nuances discussed above, the primary driver for changes in future drought remains changes in overall precipitation amount. In Texas, the projected slight decrease mentioned in the previous section would imply a small increase of drought intensity by itself. For agricultural drought, this small increase of meteorological drought could easily be neutralized by improved water use efficiency by plants, leading to no agricultural drought trend or even a reduction in drought susceptibility, though several other factors will come into play, not least the increasing temperatures. For hydrologic drought, the increase of surface water evaporation would worsen the impact of the increased meteorological drought.


Because of all the factors at play, it is impossible to make quantitative statewide projections of drought trends. The majority of factors point toward increased drought severity. Nonetheless, it seems highly likely that any such underlying trend will be dwarfed during the next couple of decades by the impact of multidecadal variability, which historical records show is large for Texas. Also, as indicated by paleoclimate records, worse droughts have occurred in Texas than the climate data record alone would indicate, and rainfall deficits comparable to those earlier in the 20th century would have greater impacts due to higher temperatures.⁵¹





RIVER FLOODING

Texas has been impacted greatly by river flooding in the past, causing both fatalities and economic damage. In fact, Texas ranked highest amongst the United States in flood related fatalities during 1959-2005.⁵² Throughout most of the state, intense daily and weekly precipitation events are the primary meteorological drivers correlated with the most extensive flooding.⁵³ This would imply that a single or a short series of extreme precipitation events plays a dominant role in causing river flooding, and that antecedent soil conditions have limited impact. However, east and northeast Texas are the exception to this pattern, as extended wet periods are the primary meteorological drivers best correlated with extensive river flooding. This would imply that antecedent soil conditions- as controlled by seasonal precipitation and evaporation- play a dominant role in flood events in those areas.

Research has found a historical decrease in the magnitude of flood events at many river gauges, resulting from abrupt drops in flood magnitude.⁵⁴ This can be attributed to the construction of reservoirs and dams for flood management throughout the 20th century. As substantial future reservoir construction is not anticipated, this trend driver will not continue. Projected increases in temperatures and precipitation intensity will have competing effects on river flooding. Increased precipitation intensity will lead to more precipitation-runoff events that suggest more river flooding in the future. In contrast, increased temperatures would mitigate flooding by decreasing soil



moisture and increasing the capacity for soil to hold new rainfall. This would limit the amount of precipitation-runoff that would go into rivers and increases in river streamflow would be lower.



Left: Locations of selected river gauges relative to river basins. **Right:** Statistical measure of 1-day 95th percentile seasonal streamflow trends of each gauge after 1965. A Mann-Kendall statistic greater than 1.96 indicates a significant increasing trend while a Mann-Kendall statistic less than -1.96 indicates a significant decreasing trend. (Adapted from Figures 1 and 14 of Mishra et al., 2011)


At the seasonal scale, there are mixed historical trends, with increased peak streamflow in the winter and decreased peak streamflow in the fall and spring that vary by climate zone.⁵⁵ For the Trinity and Neches River Basins, both in east Texas where annual precipitation is greatest within the state, most gauges show statistically significant increases in winter extreme streamflow post-1965. While the other three seasons do not have statistically significant trends, there is still a general increasing trend in extreme streamflow for the Neches and Trinity river gauges. For the Colorado and Brazos River Basins (where annual precipitation decreases towards the west), there is an increasing trend in extreme streamflow in the winter since 1965, however, the spring and fall show a general decreasing trend.

Based on limited modeling studies, it appears that the effect of increased precipitation intensity will dominate over the effect of increased temperatures, leading to an increase in peak streamflow.⁵⁶ The increase in streamflow is likely to be threshold-dependent, as soil moisture deficits will have the greatest percentage effect on runoff at low precipitation amounts. With very intense precipitation, such as with Hurricane Harvey, antecedent soil moisture will have almost no effect.⁵⁷

In summary, river flooding in Texas is projected to have no substantial change through 2036. This is in large part due to the construction of dams and reservoirs for flood management in the 20th century. There is a

mixture of historical trends categorized by season, and this does not bring forth a clear and coherent trend to project. Also, meteorological drivers of river flooding (increased rainfall intensity, decreased soil moisture) are projected to have competing influences. On balance, if an increasing trend is present in river flooding, it will be at the most extreme flood events or in the wettest parts of the state where there is so much rainfall that a decrease in soil moisture would have little mitigating impact.





URBAN FLOODING


Urban flooding differs from river flooding in that catchments tend to be much smaller and the effects of urbanization are relevant for the projection of flooding in metropolitan areas. In general, urbanization greatly decreases or even eliminates the infiltration rates of the soil through the construction of impervious surfaces. For a given precipitation event, this greatly increases the precipitation runoff within a given basin and results in higher streamflow and flooding for urban rivers. Mitigation measures, such as detention ponds, can wholly or partly counter this effect.

Historic trends of urban flooding are variable and determined by local flood control factors. The distribution of impervious surfaces within a river basin, pre-existing land surfaces, alterations of land surfaces, pre-existing flood control impoundments that alter runoff flow, the terrain within a basin, etc. all vary from city to city in Texas. Nearly all metropolitan areas in the state contain gauges with increasing, decreasing, and neutral trends in flooding, the majority of those being neutral.⁵⁸ The degree of increasing and decreasing trend gauges varies from city, with the Houston metropolitan region standing out as a hotspot for increasing urban flooding trends (with over 60% of Texas's metropolitan increasing trend gauges in Harris County alone), whereas decreasing trends are spread among many metropolitan regions. A recent study has argued that the mere existence of Houston has intensified rainfall and increased streamflow twentyfold, but those findings may not be robust.⁵⁹


Local trends in the future may not necessarily reflect what has happened in the past. Many of the decreasing trend gauges are generally the gauges with a comparatively longer record of data (over 50 years) which are usually on major rivers which have been affected by dam/reservoir construction over the record period.⁶⁰ As mentioned in the river flood section, little substantial reservoir construction is anticipated in the future. In addition, many of the increasing trend gauges tend to have a comparatively shorter record of data (25-50 years) where natural variability will have a larger influence on the calculated trend.

Regardless of the variable historical trends, Texas's urban population has increased by over 2% per year during 2000-2010, resulting in the largest urban area and the second largest urban population of the United States.⁶¹ As Texas's population continues to grow, its urban area is likely to continue to expand and become denser, and, in combination with the projected increase in intense precipitation, the effects of urbanization would result in an increase in precipitation runoff and urban flooding. Such a trend is supported by results from hydrological model simulations.⁶²

Assuming that the flooding trends in small, rapidly-responding urban basins are driven climatologically by rainfall intensity, the change in frequency of extreme rainfall would translate directly to a change in the expected frequency of urban flooding: 30%-50% more in 2036 relative to climatological expectations for 1950-1999 and 10%-15% more relative to 2000-2018.



Number of examined stream gauges exhibiting no statistically significant trends, downward trends, and upwards trends in peak river flow, grouped by metropolitan area. (Figure 4 from Berg, 2018)



Closer examination of statistically significant gauge trends in a) Dallas-Fort Worth-Arlington and b) Houston-The Woodlands-Sugar Land areas (Figure 5 from Berg, 2018)



WINTER PRECIPITATION

Winter precipitation is rare enough in Texas to be automatically disruptive in most areas. Snow and sleet are most common across northern Texas, the Panhandle, west-central Texas, and far west Texas.⁶³ However, snow has been reported on occasion in every county in Texas.

In the southern part of the state and in coastal regions, snow is rare, but nonetheless, large accumulations of snow are possible. Notable events include 1895 from Galveston to Beaumont (nearly two feet of snowfall accumulation)⁶⁴, 1985 in San Antonio (over a foot), and 2004 in Victoria (nearly a foot).⁶⁵ Given these rare events, it is perhaps not totally unbelievable that the one-day snowfall record for Texas is held not by a Panhandle city but by Hillsboro, north of Waco, with 26" in 1929. These sorts of extremely unusual but highly disruptive snow events have not been studied in a climate change context and they are too rare for any trends to be robustly detected in the historical record.

Given the size of Texas and the typical scale of snowstorms at this latitude, it is rare for individual snowstorms to affect most of the state simultaneously. Once recent such event was on February 14-16, 2021, while Texas was in the grip of a cold wave of historic proportions. That snowstorm was also historic, one of only four snowstorms in the historical record to have brought measurable snow to most of the state of Texas.

Two of the four largest snowstorms geographically were also two of the largest snowstorms in terms of maximum snowfall amount: the 1895 and 1929 snowstorms. While it's not possible to reliably estimate a trend from such a


small number of events, it is clear that such snowstorms have not increased in frequency.

Farther to the northeast, where snow is more common, **there is no noticeable trend in the frequency of 2" or greater snowfalls from 1930 to 2007.**⁶⁶ **Because the determining factor for snowfall in northwest Texas is typically air temperature, a reasonable expectation is that snowfall frequency and intensity will decrease in the future, somewhat reducing the snow hazard. In climate model projections, the risk of snowfall consistently decreases in climates like that of Texas.**⁶⁷


Freezing rain is highly disruptive to vehicular travel and to electric power transmission. Freezing rain can occur when a warm layer of air aloft overlays subfreezing air near the surface. In Texas, this happens when a strong cold front stalls across Texas as an upper-air disturbance picks up warmer, humid air from over the Gulf of Mexico and carries it northward above the shallow cold air. Freezing rain is most common in the northeastern part of the state.

The climate data record for freezing rain observations is quite sparse compared to snow and rain. Freezing rain trends in Texas during the last half of the 20th century were mixed.⁶⁸ Because freezing rain is strongly dependent on temperature conditions, one might expect a decrease in the threat of freezing rain over time. However, the weather patterns that produce freezing rain are rather specific, and changes in the frequency of those weather patterns could easily amplify or neutralize any temperature-driven trend. **So the expectation of a decline in freezing rain frequency is tentative, and the magnitude of such a decline is as yet unknown.**


% Years with Snow Reported, 1890-2021




Mean Annual Snowfall, 1890-2021



Peak Annual Snowfall, 1890-2021



Extent of Major Texas Snowstorms



Distribution of reported snowfall totals for the four largest Texas snowstorms on record. Note that stations are distributed unevenly geographically and over time; locations between reported snowfalls are likely to have received snow as well.





SEVERE THUNDERSTORMS

There is no reliable, long-term record of severe thunderstorms or the severe weather they produce: tornadoes, hail, and strong winds. Reporting methods and magnitude scales have changed over time for tornadoes and hail events.




Maps of the historic distribution of tornadoes, hail, and strong winds make it clear that no corner of the state is immune to severe thunderstorms.⁶⁹ The dense clusters of observed events around major metropolitan areas such as Dallas-Fort Worth reflect the enhanced likelihood that a severe thunderstorm there will be witnessed or will cause damage. It is also possible to detect lines of severe weather reports along some major highways, such as Interstate 20 near Midland. The strong dependence of report location on population density gives an indication of how changes in severe thunderstorm reports over time are more strongly affected by changes in population than by changes in severe thunderstorms themselves.⁷⁰

Not only is the climate data record for severe thunderstorms poor, severe thunderstorms are too

small to be simulated directly by present-day climate models. Therefore, when assessing trends in severe thunderstorms, it is necessary to consider indirect indicators of severe thunderstorm frequency and intensity such as wind shear and convective instability, both of which favor severe storms.

Over the past few decades, the severe storm environment over Texas has changed in complex and opposing ways.⁷¹ The amount of energy available for convection has decreased, and the amount of energy needed to initiate convection has increased at the same time. This suggests that environmental conditions have become less favorable for the occurrence of thunderstorms. However, the amount of low-level shear has increased, which would be expected to make thunderstorms more likely to become severe once they develop. Changes in severe storm environments have not been uniform throughout the year, with environments becoming more favorable for severe thunderstorms and significant hail in Texas early in the spring and less favorable later in the spring.

Climate model simulations imply different prospects going forward. As temperatures increase, the amount



a) Locations and magnitudes of tornadoes in Texas, 1950-2019.

b) Locations and magnitudes of severe hail in Texas, 1950-2019.

c) Locations and magnitudes of potentially damaging thunderstorm winds in Texas, 1950-2019.

Data from <https://www.spc.noaa.gov/gis/svrgis>.

of energy available to fuel these storms is simulated to increase as temperature and low-level moisture increase.⁷² Even though shear will likely decrease as the temperature gradient from the poles to the equator weakens, the increase in instability is projected to overwhelm any decrease in low level shear.⁷³ This results in an overall increase in the number of days capable of producing severe thunderstorms.

With these complex trends and partially contradictory information between models and observations, there is low confidence in any ongoing trend in the overall frequency and severity of severe thunderstorms.

Regarding the specific hazards of thunderstorms, lightning occurs most often during the months of May and June. Severe wind is most prevalent during the summer months from disorganized storm systems in the High Plains of Texas.⁷⁴ **The most robust trend in tornado activity is a tendency of more tornadoes in large outbreaks, but the factors apparently driving that trend are not projected to continue.**⁷⁵ Warmer temperatures are likely to lead to less hail overall, particular during the summer, but increases in available thunderstorm energy may lead to an increase of the risk of very large hail earlier in springtime.⁷⁶



HURRICANES AND COASTAL EROSION

Sea level rise and storm intensity both affect coastal flooding and erosion.

The change in ocean height relative to coastal lands, called relative sea level rise, is one of many factors affecting coastal erosion. Relative sea level rise is a combination of three factors: eustatic sea level rise, local variations in sea level rise, and relative land motion. Eustatic sea level rise is the change in global mean ocean height and is primarily the result of increasing temperatures that cause thermal expansion and melting glaciers and ice sheets. Local variations are produced by changes in wind patterns and ocean currents, and are minor for the Gulf of Mexico. Relative land motion in coastal Texas is dominated by coastal subsidence.

Subsidence is a gradual lowering of land-surface elevation and is the result of the extraction of groundwater, oil, or gas or increasing sediment loading or infrastructure construction. As the coast of Texas slowly sinks, water potentially encroaches landward so quickly that it can exceed natural sediment accretion rates. In the state of Texas, the rate of subsidence ranges from less than 2 mm/yr to as much as 7 mm/yr.⁷⁸ The variations are due to historical differences in oil, gas, or groundwater extraction and sediment loading, resulting in generally larger rates of subsidence in southeast Texas than in south Texas.⁷⁹

The combination of local subsidence, eustatic sea level rise, and changes in sediment deposition and transport have produced a retreat of the Texas coastline along nearly the entire length of its barrier islands.⁸⁰ In

	Station ID		RSLR	95% C
Sabine Pass	8770822	1958-2020	6.16 mm/yr	+/- 0.74
Galveston Pier 21	8771450	1904-2020	6.59 mm/yr	+/- 0.22
Freeport	8772440	1972-2008	4.43 mm/yr	+/- 1.05
Rockport	8774770	1937-2020	5.86 mm/yr	+/- 0.48
Corpus Christi	8775870	1983-2020	5.44 mm/yr	+/- 1.04
Port Mansfield	8778490	1963-2020	3.54 mm/yr	+/- 0.70
Port Isabel	8779770	1944-2020	4.18 mm/yr	+/- 0.30
S. Padre Island	8779748	1958-2020	4.27 mm/yr	+/- 0.58

Relative sea-level rise (RSLR) and 95% confidence interval (95% C) at selected Texas tide gauges through 2020.⁷⁷


Galveston Bay and probably other bays and estuaries behind the barrier islands, sediment deposition is not keeping up with relative sea level rise, leading to loss of coastal wetlands.⁸¹ In many areas there has been a decrease in extraction of groundwater and other resources as the problems associated with coastal subsidence have become clearer, resulting in a reduction in coastal subsidence. However, eustatic sea level has shown indications of acceleration,⁸² so on balance the near-term future rate of coastline retreat may be expected to be similar to historic rates.

Rising sea levels lead directly to increased risk of storm surge from hurricanes, as the storm surge is on top of an elevated baseline. Given typical return periods for storm surges along the Gulf Coast,⁸³ a 1 meter relative sea level rise produces a doubling of storm surge risk, as a surge that would in the past have been expected have a 1% chance of occurring in any given year would in the future have a 2% chance of occurring in any given year. **The places along the coast with the largest rates of relative**

sea level rise may have a doubled storm surge risk by 2050 relative to the risk at the beginning of the 20th century, purely due to the relative sea level rise itself.

An additional element of enhanced risk is provided by an expected increase in the intensity of very strong hurricanes, despite an expected lack of increase, or even a decrease, in hurricane frequency overall.⁸⁴ Different research studies have produced some conflicting results, and local trends over the western Gulf of Mexico will also be affected by changes in wind patterns for which global climate models have little predictive skill. While some recent research has pointed to an apparent trend for United States tropical cyclones to move more slowly at landfall, much like Hurricane Harvey, other research suggests that Texas may be spared from such a slowdown.⁸⁵ At this point, the enhanced risk is difficult to quantify, but substantial scientific progress on this topic is likely as climate models become better able to simulate the observed spatial distribution, frequency, and intensity of hurricanes.⁸⁶

Net rates of coastal advance (positive) or retreat (negative) for the Texas Gulf shoreline between Sabine Pass and the Rio Grande, calculated from shoreline positions between 2000 and 2019. Longer-term changes are similar. From Paine and Caudle (2020).





WILDFIRES

Fire depends on strong winds, low humidity, surface dryness, and factors affecting the amount of vegetation available to burn (known as fuel load).

There is weak statistical evidence for a slight decline in the frequency of unusually strong winds speeds across the southern Great Plains.⁸⁷ One future climate simulation predicts an increase in average wind speed across most of Texas in all seasons.⁸⁸ On the whole, expectations for wind speed trends lack robustness.

Low humidity and surface dryness are closely related. Low humidity refers to the extremely dry weather accompanying wildfire outbreaks across Texas, while surface dryness refers to the lack of moisture in dry, dormant, or otherwise combustible vegetation. Surface dryness is strongly influenced by weather conditions, but it is a consequence of desiccation of vegetation over an extended period of time rather than simply lack of humidity on a particular day.

Vapor pressure deficit, an indicator of the ability of moisture to evaporate, is projected to increase as temperatures rise and carbon dioxide fertilization reduces transpiration, leading to both lower humidity and increased surface dryness.⁸⁹ **Overall, increased dryness should extend the wildfire season in places where the fire season is presently constrained by low levels of aridity, such as eastern Texas.**⁹⁰

Changes in fuel load involve two competing effects: increased aridity leading to reduced plant growth, and increased carbon dioxide leading to increased plant growth. Multiple papers predict wildfires in the Southwest (including Texas) will change differently than elsewhere in the country as fuel load gradually becomes the determining limit on fires.⁹¹ **This transition in wildfire regime suggests that over the next twenty years, wildfire risk may increase more slowly in the Panhandle and Far West Texas than elsewhere as increased aridity reduces biomass. Meanwhile, the area of the state commonly affected by wildfires may expand eastward as fuels become drier faster in a warmer climate.**

REFERENCES

- Al Mukaimi, M. E., T. M. Dellapenna, and J. R. Williams, 2018: Enhanced land subsidence in Galveston Bay, Texas: Interaction between sediment accumulation rates and relative sea level rise. *Estuarine, Coastal and Shelf Science*, **207**, 183-193.
- Anderson, C. J., C. K. Wikle, Q. Zhou, and J. A. Royle, 2007: Population Influences on Tornado Reports in the United States. *Weather and Forecasting*, **22**, 571-579.
- Ashley, S. T., and W. S. Ashley, 2008: Flood Fatalities in the United States. *Journal of Applied Meteorology and Climatology*, **47**, 805-818.
- Berg, A., J. Sheffield, and P. C. D. Milly, 2017: Divergent surface and total soil moisture projections under global warming. *Geophysical Research Letters*, **44**, 236-244.
- Berg, A., and Coauthors, 2016: Land-atmosphere feedbacks amplify aridity increase over land under global warming. *Nature Climate Change*, **6**, 869-+.
- Berg, M. D., 2018: Peak flow trends highlight emerging urban flooding hotspots in Texas. *Texas Water Journal*, **9**, 18-29.
- Berghuijs, W. R., R. A. Woods, C. J. Hutton, and M. Sivapalan, 2016: Dominant flood generating mechanisms across the United States. *Geophysical Research Letters*, **43**, 4382-4390.
- Blackport, R., J. A. Screen, K. van der Wiel, and R. Bintanja, 2019: Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes. *Nature Climate Change*, **9**, 697-704.
- Bomar, G. W., 2017: *Weather in Texas: The Essential Handbook*. Third Edition ed. University of Texas Press, 290 pp.
- Brimelow, J. C., W. R. Burrows, and J. M. Hanesiak, 2017: The changing hail threat over North America in response to anthropogenic climate change. *Nature Climate Change*, **7**, 516.
- Brooks, H. E., 2013: Severe thunderstorms and climate change. *Atmospheric Research*, **123**, 129-138.
- Bukovsky, M. S., R. R. McCrary, A. Seth, and L. O. Mearns, 2017: A Mechanistically Credible, Poleward Shift in Warm-Season Precipitation Projected for the U.S. Southern Great Plains? *Journal of Climate*, **30**, 8275-8298.
- Camargo, S. J., and A. A. Wing, 2016: Tropical cyclones in climate models. *Wiley Interdisciplinary Reviews: Climate Change*, **7**, 211-237.
- Capotondi, A., and M. A. Alexander, 2010: Relationship between Precipitation in the Great Plains of the United States and Global SSTs: Insights from the IPCC AR4 Models. *Journal of Climate*, **23**, 2941-2958.
- Capotondi, A., and Coauthors, 2015: Understanding ENSO Diversity. *Bulletin of the American Meteorological Society*, **96**, 921-938.
- Changnon, S. A., and T. R. Karl, 2003: Temporal and Spatial Variations of Freezing Rain in the Contiguous United States: 1948-2000. *Journal of Applied Meteorology*, **42**, 1302-1315.
- Cleveland, M. K., T. H. Votteler, D. K. Stahle, R. C. Casteel, and J. L. Banner, 2011: Extended Chronology of Drought in South Central, Southeastern and West Texas. *Texas Water Journal*, **2**, 54-96.
- Cohen, J., K. Pfeiffer, and J. A. Francis, 2018: Warm Arctic episodes linked with increased frequency of extreme winter weather in the United States. *Nature Communications*, **9**, 869.
- Cohen, J., L. Agel, M. Barlow, C. I. Garfinkel, and I. White, 2021: Linking Arctic variability and change with extreme winter weather in the United States. *Science*, **373**, 1116-1121.
- Cohen, J., and Coauthors, 2014: Recent Arctic amplification and extreme mid-latitude weather. *Nature Geoscience*, **7**, 627.
- Collins, M., and Coauthors, 2013: Long-term Climate Change: Projections, Commitments and Irreversibility. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker, and Coauthors, Eds., Cambridge University Press.
- Cook, B. I., T. R. Ault, and J. E. Smerdon, 2015: Unprecedented 21st century drought risk in the American Southwest and Central Plains. *Science Advances*.
- Dai, A., T. Zhao, and J. Chen, 2018: Climate Change and Drought: a Precipitation and Evaporation Perspective. *Current Climate Change Reports*, **4**, 301-312.

- Dangendorf, S., C. Hay, F. M. Calafat, M. Marcos, C. G. Piecuch, K. Berk, and J. Jensen, 2019: Persistent acceleration in global sea-level rise since the 1960s. *Nature Climate Change*, **9**, 705-710.
- Davis, R. A. J., 2011: *Sea-Level Change in the Gulf of Mexico*. Texas A&M University Press, 171 pp.
- Diffenbaugh, N. S., M. Scherer, and R. J. Trapp, 2013: Robust increases in severe thunderstorm environments in response to greenhouse forcing. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, 16361-16366.
- Easterling, D. R., and Coauthors, 2017: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 207-230.
- Elsner, J. B., L. E. Michaels, K. N. Scheitlin, and I. J. Elsner, 2013: The Decreasing Population Bias in Tornado Reports across the Central Plains. *Weather, Climate, and Society*, **5**, 221-232.
- Frederikse, T., S. Jevrejeva, R. E. M. Riva, and S. Dangendorf, 2018: A Consistent Sea-Level Reconstruction and Its Budget on Basin and Global Scales over 1958-2014. *Journal of Climate*, **31**, 1267-1280.
- Guyette, R. P., F. R. Thompson, J. Whittier, M. C. Stambaugh, and D. C. Dey, 2014: Future Fire Probability Modeling with Climate Change Data and Physical Chemistry. *Forest Science*, **60**, 862-870.
- Hall, T. M., and J. P. Kossin, 2019: Hurricane stalling along the North American coast and implications for rainfall. *npj Climate and Atmospheric Science*, **2**, 17.
- Hassanzadeh, P., C.-Y. Lee, E. Nabizadeh, S. J. Camargo, D. Ma, and L. Y. Yeung, 2020: Effects of climate change on the movement of future landfalling Texas tropical cyclones. *Nature Communications*, **11**, 3319.
- Huang, Y., S. Wu, and J. O. Kaplan, 2015: Sensitivity of global wildfire occurrences to various factors in the context of global change. *Atmospheric Environment*, **121**, 86-92.
- Iturbide, M., and Coauthors, 2021: Repository supporting the implementation of FAIR principles in the IPCC-WG1 Atlas.
- Janssen, E., D. J. Wuebbles, K. E. Kunkel, S. C. Olsen, and A. Goodman, 2014: Observational- and model-based trends and projections of extreme precipitation over the contiguous United States. *Earths Future*, **2**, 99-113.
- Kirtman, B., and Coauthors, 2013: Near-term Climate Change: Projections and Predictability. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker, and Coauthors, Eds., Cambridge University Press.
- Kloesel, K., and Coauthors, 2018: Southern Great Plains. *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II*, D. R. Reidmiller, C. W. Avery, D. R. Easterling, K. E. Kunkel, K. L. M. Lewis, T. K. Maycock, and B. C. Stewart, Eds., U.S. Global Change Research Program, 987-1035.
- Kliver, D., and D. Leathers, 2015: Winter snowfall prediction in the United States using multiple discriminant analysis. *International Journal of Climatology*, **35**, 2003-2018.
- Knutson, T., and Coauthors, 2019: Tropical Cyclones and Climate Change Assessment: Part I: Detection and Attribution. *Bulletin of the American Meteorological Society*, **100**, 1987-2007.
- , 2020: Tropical Cyclones and Climate Change Assessment: Part II: Projected Response to Anthropogenic Warming. *Bulletin of the American Meteorological Society*, **101**, E303-E322.
- Koch, E., J. Koh, A. C. Davison, C. Lepore, and M. K. Tippett, 2021: Trends in the Extremes of Environments Associated with Severe U.S. Thunderstorms. *Journal of Climate*, **34**, 1259-1272.
- Kossin, J. P., T. Hall, T. R. Knutson, K. E. Kunkel, R. J. Trapp, D. E. Waliser, and M. E. Wehner, 2017: Extreme Storms. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 257-276.
- Kunkel, K. E., and Coauthors, 2013: Monitoring and Understanding Trends in Extreme Storms: State of Knowledge. *Bulletin of the American Meteorological Society*, **94**, 499-514.
- Kurtzman, D., and B. R. Scanlon, 2007: El Nino-Southern Oscillation and Pacific Decadal Oscillation impacts on precipitation in the southern and central United States: Evaluation of spatial distribution and predictions. *Water Resources Research*, **43**.
- Kushnir, Y., R. Seager, M. Ting, N. Naik, and J. Nakamura, 2010: Mechanisms of Tropical Atlantic SST Influence on North American Precipitation Variability*. *Journal of Climate*, **23**, 5610-5628.

- Lanza, M., 2017: Space City Rewind: Houston's Great Snow of 1895. *Space City Weather*.
- Letetrel, C., M. Karpytchev, M. N. Bouin, M. Marcos, A. Santamaría-Gómez, and G. Wöppelmann, 2015: Estimation of vertical land movement rates along the coasts of the Gulf of Mexico over the past decades. *Continental Shelf Research*, **111**, 42-51.
- Li, X., and Coauthors, 2020: Impacts of urbanization, antecedent rainfall event, and cyclone tracks on extreme floods at Houston reservoirs during Hurricane Harvey. *Environmental Research Letters*, **15**, 124012.
- Liu, Y., S. L. Goodrick, and J. A. Stanturf, 2013: Future U.S. wildfire potential trends projected using a dynamically downscaled climate change scenario. *Forest Ecology and Management*, **294**, 120-135.
- Maloney, E. D., and Coauthors, 2014: North American Climate in CMIP5 Experiments: Part III: Assessment of Twenty-First-Century Projections. *Journal of Climate*, **27**, 2230-2270.
- McCabe, G. J., M. A. Palecki, and J. L. Betancourt, 2004: Pacific and Atlantic Ocean influences on multidecadal drought frequency in the United States. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 4136-4141.
- Meehl, G. A., J. M. Arblaster, and C. T. Y. Chung, 2015: Disappearance of the southeast US "warming hole" with the late 1990s transition of the Interdecadal Pacific Oscillation. *Geophysical Research Letters*, **42**, 5564-5570.
- Milly, P. C. D., and K. A. Dunne, 2016: Potential evapotranspiration and continental drying. *Nature Climate Change*, **6**, 946-+.
- Mishra, A. K., and V. P. Singh, 2010: Changes in extreme precipitation in Texas. *Journal of Geophysical Research-Atmospheres*, **115**.
- Mishra, A. K., V. P. Singh, and M. Özger, 2011: Seasonal streamflow extremes in Texas river basins: Uncertainty, trends, and teleconnections. *Journal of Geophysical Research*, **116**.
- Morice, C. P., and Coauthors, 2021: An Updated Assessment of Near-Surface Temperature Change From 1850: The HadCRUT5 Data Set. *Journal of Geophysical Research: Atmospheres*, **126**, e2019JD032361.
- Mukherjee, S., A. Mishra, and K. E. Trenberth, 2018: Climate Change and Drought: a Perspective on Drought Indices. *Current Climate Change Reports*, **4**, 145-163.
- Needham, H. F., B. D. Keim, D. Sathiaraj, and M. Shafer, 2012: Storm Surge Return Periods for the United States Gulf Coast. *Advances in Hurricane Engineering*, 715-740.
- Nerem, R. S., B. D. Beckley, J. T. Fasullo, B. D. Hamlington, D. Masters, and G. T. Mitchum, 2018: Climate-change-driven accelerated sea-level rise detected in the altimeter era. *Proceedings of the National Academy of Sciences of the United States of America*, **115**, 2022-2025.
- Nielsen-Gammon, J. W., and Coauthors, 2020: Unprecedented Drought Challenges for Texas Water Resources in a Changing Climate: What Do Researchers and Stakeholders Need to Know? *Earth's Future*, **8**, e2020EF001552.
- Nigam, S., B. Guan, and A. Ruiz-Barradas, 2011: Key role of the Atlantic Multidecadal Oscillation in 20th century drought and wet periods over the Great Plains. *Geophysical Research Letters*, **38**, n/a-n/a.
- Nikiel, C. A., and E. A. B. Eltahir, 2019: Summer Climate Change in the Midwest and Great Plains due to Agricultural Development during the Twentieth Century. *Journal of Climate*, **32**, 5583-5599.
- O'Gorman, P. A., 2014: Contrasting responses of mean and extreme snowfall to climate change. *Nature*, **512**, 416-418.
- Overland, J. E., and Coauthors, 2021: How do intermittency and simultaneous processes obfuscate the Arctic influence on midlatitude winter extreme weather events? *Environmental Research Letters*, **16**, 043002.
- Paciorek, C. J., D. A. Stone, and M. F. Wehner, 2018: Quantifying statistical uncertainty in the attribution of human influence on severe weather. *Weather and Climate Extremes*, **20**, 69-80.
- Paine, J. G., and T. L. Caudle, 2020: Shoreline Movement along the Texas Gulf Coast, 1930s to 2019, 64 pp.
- Paine, J. G., T. L. Caudle, and J. R. Andrews, 2017: Shoreline and Sand Storage Dynamics from Annual Airborne LIDAR Surveys, Texas Gulf Coast. *Journal of Coastal Research*, **33**, 487-506.
- Pendergrass, A. G., R. Knutti, F. Lehner, C. Deser, and B. M. Sanderson, 2017: Precipitation variability increases in a warmer climate. *Sci Rep*, **7**, 17966.
- Perica, S., S. Pavlovic, M. St. Laurent, C. Trypaluk, D. Unruh, and O. Wilhite, 2018: NOAA Atlas 14: Precipitation-Frequency Atlas of the United States, Volume 11 Version 2.0: Texas, 40 pp.

- Pfahl, S., P. A. O’Gorman, and E. M. Fischer, 2017: Understanding the regional pattern of projected future changes in extreme precipitation. *Nature Climate Change*, **7**, 423.
- Pryor, S. C., and J. Ledolter, 2010: Addendum to “Wind speed trends over the contiguous United States”. *Journal of Geophysical Research: Atmospheres*, **115**.
- Qing, Y., S. Wang, B. Zhang, and Y. Wang, 2020: Ultra-high resolution regional climate projections for assessing changes in hydrological extremes and underlying uncertainties. *Climate Dynamics*, **55**, 2031-2051.
- Scheff, J., 2018: Drought Indices, Drought Impacts, CO₂, and Warming: a Historical and Geologic Perspective. *Current Climate Change Reports*, **4**, 202-209.
- Schmandt, J., G. R. North, and J. Clarkson, 2011: *The Impact of Global Warming on Texas*. Second ed. University of Texas Press, 318 pp.
- Schubert, S., and Coauthors, 2009: A U.S. CLIVAR Project to Assess and Compare the Responses of Global Climate Models to Drought-Related SST Forcing Patterns: Overview and Results. *Journal of Climate*, **22**, 5251-5272.
- Seager, R., A. Hooks, A. P. Williams, B. Cook, J. Nakamura, and N. Henderson, 2015: Climatology, Variability, and Trends in the U.S. Vapor Pressure Deficit, an Important Fire-Related Meteorological Quantity. *Journal of Applied Meteorology and Climatology*, **54**, 1121-1141.
- Smith, B. T., T. E. Castellanos, A. C. Winters, C. M. Mead, A. R. Dean, and R. L. Thompson, 2013: Measured Severe Convective Wind Climatology and Associated Convective Modes of Thunderstorms in the Contiguous United States, 2003–09. *Weather and Forecasting*, **28**, 229-236.
- Stambaugh, M. C., R. P. Guyette, E. D. Stroh, M. A. Struckhoff, and J. B. Whittier, 2018: Future southcentral US wildfire probability due to climate change. *Climatic Change*, **147**, 617-631.
- Swann, A. L. S., 2018: Plants and Drought in a Changing Climate. *Current Climate Change Reports*, **4**, 192-201.
- Swann, A. L. S., F. M. Hoffman, C. D. Koven, and J. T. Randerson, 2016: Plant responses to increasing CO₂ reduce estimates of climate impacts on drought severity. *Proceedings of the National Academy of Sciences of the United States of America*, **113**, 10019-10024.
- Tang, B. H., V. A. Gensini, and C. R. Homeyer, 2019: Trends in United States large hail environments and observations. *npj Climate and Atmospheric Science*, **2**, 45.
- Taszarek, M., J. T. Allen, H. E. Brooks, N. Pilgaj, and B. Czernecki, 2021: Differing Trends in United States and European Severe Thunderstorm Environments in a Warming Climate. *Bulletin of the American Meteorological Society*, **102**, E296-E322.
- Thomson, A. M., and Coauthors, 2011: RCP4.5: a pathway for stabilization of radiative forcing by 2100. *Climatic Change*, **109**, 77-94.
- Tippett, M. K., C. Lepore, and J. E. Cohen, 2016: More tornadoes in the most extreme U.S. tornado outbreaks. *Science*, **354**, 1419-1423.
- Trenberth, K. E., A. Dai, R. M. Rasmussen, and D. B. Parsons, 2003: The Changing Character of Precipitation. *Bulletin of the American Meteorological Society*, **84**, 1205-1218.
- TSHA, 2018: *Texas Almanac 2018-2019*. Texas State Historical Association, 752 pp.
- USGCRP, 2017: *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. U.S. Global Change Research Program, 470 pp.
- Van Klooster, S. L., and P. J. Roebber, 2009: Surface-Based Convective Potential in the Contiguous United States in a Business-as-Usual Future Climate. *Journal of Climate*, **22**, 3317-3330.
- van Oldenborgh, G. J., E. Mitchell-Larson, G. A. Vecchi, H. de Vries, R. Vautard, and F. Otto, 2019: Cold waves are getting milder in the northern midlatitudes. *Environmental Research Letters*, **14**, 114004.
- van Vuuren, D. P., and Coauthors, 2011: The representative concentration pathways: an overview. *Climatic Change*, **109**, 5-31.
- Villarini, G., and J. A. Smith, 2013: FLOODING IN TEXAS: EXAMINATION OF TEMPORAL CHANGES AND IMPACTS OF TROPICAL CYCLONES. *Journal of the American Water Resources Association*, **49**, 825-837.
- Vose, R. S., D. R. Easterling, K. E. Kunkel, A. N. LeGrande, and M. F. Wehner, 2017: Temperature changes in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 185-206.
- Vose, R. S., and Coauthors, 2014: Improved historical temperature and precipitation time series for U.S. climate divisions. *Journal of Applied Meteorology and Climatology*, **53**, 1232-1251.

Wang, H., S. Schubert, M. Suarez, and R. Koster, 2010: The Physical Mechanisms by Which the Leading Patterns of SST Variability Impact U.S. Precipitation. *Journal of Climate*, **23**, 1815-1836.

Wood, A. W., L. R. Leung, V. Sridhar, and D. P. Lettenmaier, 2004: Hydrologic implications of dynamical and statistical approaches to downscaling climate model outputs. *Climatic Change*, **62**, 189-216.

Zhang, W., G. Villarini, G. A. Vecchi, and J. A. Smith, 2018: Urbanization exacerbated the rainfall and flooding caused by hurricane Harvey in Houston. *Nature*, **563**, 384-388.

Zhao, G., H. L. Gao, and L. Cuo, 2016: Effects of Urbanization and Climate Change on Peak Flows over the San Antonio River Basin, Texas. *Journal of Hydrometeorology*, **17**, 2371-2389.



ENDNOTES

ⁱ More general discussions of climate change and its impact on Texas may be found in: Schmandt, J., G. R. North, and J. Clarkson, 2011: *The Impact of Global Warming on Texas*. Second ed. University of Texas Press, 318 pp.

USGCRP, 2017: *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. U.S. Global Change Research Program, 470 pp.

Kloesel, K., and Coauthors, 2018: Southern Great Plains. *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II*, D. R. Reidmiller, C. W. Avery, D. R. Easterling, K. E. Kunkel, K. L. M. Lewis, T. K. Maycock, and B. C. Stewart, Eds., U.S. Global Change Research Program, 987-1035.

ⁱⁱ Online data archive: <ftp://ftp.ncei.noaa.gov/pub/data/cirs/climdiv/>

Vose, R. S., and Coauthors, 2014: Improved historical temperature and precipitation time series for U.S. climate divisions. *Journal of Applied Meteorology and Climatology*, **53**, 1232-1251.

ⁱⁱⁱ Global temperatures are from the HadCRUT4 data set. All linear trends in this report are calculated using ordinary least squares. Morice, C. P., and Coauthors, 2021: An Updated Assessment of Near-Surface Temperature Change From 1850: The HadCRUT5 Data Set. *Journal of Geophysical Research: Atmospheres*, **126**, e2019JD032361.

^{iv} Meehl, G. A., J. M. Arblaster, and C. T. Y. Chung, 2015: Disappearance of the southeast US "warming hole" with the late 1990s transition of the Interdecadal Pacific Oscillation. *Geophysical Research Letters*, **42**, 5564-5570, Vose, R. S., D. R. Easterling, K. E. Kunkel, A. N. LeGrande, and M. F. Wehner, 2017: Temperature changes in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 185-206.

^v The historical simulations and projections are based on CMIP5 multi-model ensemble output from the Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections archive at https://gdo-dcp.ucllnl.org/downscaled_cmip_projections

The first available ensemble member from the combined Historic and RCP 4.5 was used from each of the following models for temperature and from the combined Historic and RCP 8.5 for precipitation. A ^ means only RCP 4.5 simulations were available from a particular model, so it was not included in the precipitation ensemble.

Modeling Center (or Group)	Institute ID	Model Name
Commonwealth Scientific and Industrial Research Organization (CSIRO) and Bureau of Meteorology (BOM), Australia	CSIRO-BOM	ACCESS1.0 ACCESS1.3 [^]
Beijing Climate Center, China Meteorological Administration	BCC	BCC-CSM1.1 BCC-CSM1.1(m)
Canadian Centre for Climate Modelling and Analysis	CCCMA	CanESM2
National Center for Atmospheric Research	NCAR	CCSM4
Community Earth System Model Contributors	NSF-DOE-NCAR	CESM1(BGC) CESM1(CAM5)
Centro Euro-Mediterraneo per I Cambiamenti Climatici	CMCC	CMCC-CM
Centre National de Recherches Météorologiques / Centre Européen de Recherche et Formation Avancée en Calcul Scientifique	CNRM-CERFACS	CNRM-CM5
Commonwealth Scientific and Industrial Research Organization in collaboration with Queensland Climate Change Centre of Excellence	CSIRO-QCCCE	CSIRO-Mk3.6.0

EC-EARTH consortium	EC-EARTH	EC-EARTH [^]
LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences and CESS, Tsinghua University	LASG-CESS	FGOALS-g2
LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences	LASG-IAP	FGOALS-s2 [^]
The First Institute of Oceanography, SOA, China	FIO	FIO-ESM
NOAA Geophysical Fluid Dynamics Laboratory	NOAA GFDL	GFDL-CM3 GFDL-ESM2G GFDL-ESM2M
NASA Goddard Institute for Space Studies	NASA GISS	GISS-E2-H-CC [^] GISS-E2-R GISS-E2-R-CC [^]
National Institute of Meteorological Research/Korea Meteorological Administration	NIMR/KMA	HadGEM2-AO
Met Office Hadley Centre (additional HadGEM2-ES realizations contributed by Instituto Nacional de Pesquisas Espaciais)	MOHC (additional realizations by INPE)	HadGEM2-CC HadGEM2-ES
Institute for Numerical Mathematics	INM	INM-CM4
Institut Pierre-Simon Laplace	IPSL	IPSL-CM5A-LR [^] IPSL-CM5A-MR IPSL-CM5B-LR
Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies	MIROC	MIROC-ESM MIROC-ESM-CHEM
Atmosphere and Ocean Research Institute (The University of Tokyo), National Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology	MIROC	MIROC5
Max-Planck-Institut für Meteorologie (Max Planck Institute for Meteorology)	MPI-M	MPI-ESM-MR MPI-ESM-LR
Meteorological Research Institute	MRI	MRI-CGCM3
Norwegian Climate Centre	NCC	NorESM1-M NorESM1-ME [^]

The 1/8-degree BCSD (bias-corrected and statistically-downscaled) model output was downloaded as a geographically averaged ensemble mean over a latitude-longitude box centered on Texas: 27.5°-35.5°N, 94.25°-104.0°W, inclusive.

Wood, A. W., L. R. Leung, V. Sridhar, and D. P. Lettenmaier, 2004: Hydrologic implications of dynamical and statistical approaches to downscaling climate model outputs. *Climatic Change*, **62**, 189-216.

^{vi} van Vuuren, D. P., and Coauthors, 2011: The representative concentration pathways: an overview. *ibid.*, **109**, 5-31. Thomson, A. M., and Coauthors. *ibid.*: RCP4.5: a pathway for stabilization of radiative forcing by 2100, 77-94.

^{vii} Kirtman, B., and Coauthors, 2013: Near-term Climate Change: Projections and Predictability. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker, and Coauthors, Eds., Cambridge University Press.

^{viii} 1900-1999 average: 64.6 °F. 2000-2018 average: 66.0 °F. 2036 projection: 67.6 °F. Warmest year on record: 2012 (67.8 °F).

^{ix} The stations are chosen for their geographical representativeness, data completeness, and lack of substantial changes in station location. These stations, and which of four regions of the state in which they are found (Northwest, Northeast, South, and Coastal), are listed here:

Urban: El Paso Intl Airport (NW); San Antonio Intl Airport (S); Dallas Love Field (NE); Houston Hobby Airport (C).

Semi-Urban: Hereford (NW); Huntsville (NE); College Station Easterwood Field (S); Harlingen (C); Corpus Christi Intl Airport (C); Brownfield (NW)

Rural: Dalhart FAA Airport (NW); Luling (S); Midland Intl Airport (NW); Lufkin Angelina Co Airport (NE); Amarillo Intl Airport (NW); Victoria Regional Airport (C); Beeville 5 NE (R); Crosbyton (NW); Centerville (NE)
In order to retain a sufficient number of homogeneous stations, the index data record only covers the period 1950-present.

^x The county temperature stations are created as follows: using SC-ACIS web services, every daily maximum/minimum temperature observation is retrieved for a given county. Each year, starting in 1890, a station with no more than ten missing days is identified. If no such station exists, data for that year is marked as missing. If at least one such station exists, data from the station with the longest period-of-record is chosen. This data becomes the year's data for that county. The process is repeated for each year through June 2021. Compared to the index stations, the county stations are much less homogeneous. In particular, different stations would have taken observations at different times, which can skew average temperatures. Because of this, we only use the county stations to identify trends in monthly temperature extremes. Such block maxima are relatively insensitive to time-of-observation artifacts.

^{xi} A reference temperature was identified for each station, corresponding to the hottest 5% of summer days. Temperature probabilities and trends were collectively analyzed for all stations within a region relative to the reference temperatures. Finally, the regional temperature distributions and trends were applied to the temperatures of a typical station within each region to assess changes in 100 °F days. Because 100 °F day counts are a nonlinear function of temperature and because counts cannot be negative, trends were analyzed on the basis of a linear fit to the time series of the logarithm of annual 100 °F day counts.

^{xii} Each year's value consists of the average of four temperatures: the hottest temperature in June, the hottest temperature in July, the hottest temperature in August, and the hottest temperature in September. For example, if at the station being used for Travis County the highest temperature during June 2008 was 95, in July 2008 98, in August 2008 100, and in September 2008 97, the average hottest day in 2008 would be 97.5.

^{xiii} USGCRP, 2017: *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. U.S. Global Change Research Program, 470 pp.

^{xiv} Nikiel, C. A., and E. A. B. Eltahir, 2019: Summer Climate Change in the Midwest and Great Plains due to Agricultural Development during the Twentieth Century. *Journal of Climate*, **32**, 5583-5599.

^{xv} Calculated from county stations as the average of the lowest temperatures recorded during each of December, January, and February, and assigned to January's year.

^{xvi} Cohen, J., and Coauthors, 2014: Recent Arctic amplification and extreme mid-latitude weather. *Nature Geoscience*, **7**, 627.

Cohen, J., K. Pfeiffer, and J. A. Francis, 2018: Warm Arctic episodes linked with increased frequency of extreme winter weather in the United States. *Nature Communications*, **9**, 869.

Blackport, R., J. A. Screen, K. van der Wiel, and R. Bintanja, 2019: Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes. *Nature Climate Change*, **9**, 697-704.

Overland, J. E., and Coauthors, 2021: How do intermittency and simultaneous processes obfuscate the Arctic influence on midlatitude winter extreme weather events? *Environmental Research Letters*, **16**, 043002.

Cohen, J., L. Agel, M. Barlow, C. I. Garfinkel, and I. White, 2021: Linking Arctic variability and change with extreme winter weather in the United States. *Science*, **373**, 1116-1121.

^{xvii} van Oldenborgh, G. J., E. Mitchell-Larson, G. A. Vecchi, H. de Vries, R. Vautard, and F. Otto, 2019: Cold waves are getting milder in the northern midlatitudes. *Environmental Research Letters*, **14**, 114004.

^{xviii} The county precipitation stations were created following the same method as the county temperature stations.

^{xix} Precipitation from single ensemble members from the Historical+RCP4.5 CMIP5 runs were averaged over the box 25°N-37.5°N, 95°W-105°W and downloaded from the KNMI Climate Explorer. The models were: ACCESS1-0 ACCESS1-3 bcc-csm1-1 bcc-csm1-1-m BNU-ESM CanESM2 CCSM4 CESM1-BGC CESM1-CAM5 CMCC-CM CMCC-CMS CNRM-CM5 CSIRO-Mk3-6-0 EC-EARTH FGOALS-g2 FIO-ESM GFDL-CM3 GFDL-ESM2G GFDL-ESM2M GISS-E2-H GISS-E2-H GISS-E2-H GISS-E2-H-CC GISS-E2-R GISS-E2-R GISS-E2-R GISS-E2-R-CC HadGEM2-AO HadGEM2-CC HadGEM2-ES inmcm4 IPSL-CM5A-LR IPSL-CM5A-MR IPSL-CM5B-LR

MIROC5 MIROC-ESM MIROC-ESM-CHEM MPI-ESM-LR MPI-ESM-MR MRI-CGCM3 NorESM1-M NorESM1-ME.

^{xx} Maloney, E. D., and Coauthors, 2014: North American Climate in CMIP5 Experiments: Part III: Assessment of Twenty-First-Century Projections. *Journal of Climate*, **27**, 2230-2270.

^{xxi} McCabe, G. J., M. A. Palecki, and J. L. Betancourt, 2004: Pacific and Atlantic Ocean influences on multidecadal drought frequency in the United States. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 4136-4141.

Kurtzman, D., and B. R. Scanlon, 2007: El Niño-Southern Oscillation and Pacific Decadal Oscillation impacts on precipitation in the southern and central United States: Evaluation of spatial distribution and predictions. *Water Resources Research*, **43**.

Schubert, S., and Coauthors, 2009: A U.S. CLIVAR Project to Assess and Compare the Responses of Global Climate Models to Drought-Related SST Forcing Patterns: Overview and Results. *Journal of Climate*, **22**, 5251-5272.

Kushnir, Y., R. Seager, M. Ting, N. Naik, and J. Nakamura, 2010: Mechanisms of Tropical Atlantic SST Influence on North American Precipitation Variability**ibid.*, **23**, 5610-5628.

Wang, H., S. Schubert, M. Suarez, and R. Koster*ibid.*: The Physical Mechanisms by Which the Leading Patterns of SST Variability Impact U.S. Precipitation, 1815-1836.

Capotondi, A., and M. A. Alexander*ibid.*: Relationship between Precipitation in the Great Plains of the United States and Global SSTs: Insights from the IPCC AR4 Models, 2941-2958.

Nigam, S., B. Guan, and A. Ruiz-Barradas, 2011: Key role of the Atlantic Multidecadal Oscillation in 20th century drought and wet periods over the Great Plains. *Geophysical Research Letters*, **38**, n/a-n/a.

^{xxii} Texas precipitation data is from the nClimDiv dataset of the National Centers for Environmental Information. The AMO and PDO were obtained from the KNMI Climate Explorer using ERSSTv5 sea surface temperatures and, in the case of the AMO, the AMO residual from a regression against global mean surface temperature.

^{xxiii} Bukovsky et al. (2017) argued for a plausible tendency for less precipitation in Texas in the summertime in CMIP5 models, but CMIP6 model simulations do not have a clear summertime drying signal in Texas.

Bukovsky, M. S., R. R. McCrary, A. Seth, and L. O. Mearns, 2017: A Mechanistically Credible, Poleward Shift in Warm-Season Precipitation Projected for the U.S. Southern Great Plains? *Journal of Climate*, **30**, 8275-8298.

Iturbide, M., and Coauthors, 2021: Repository supporting the implementation of FAIR principles in the IPCC-WG1 Atlas.

^{xxiv} Easterling, D. R., and Coauthors, 2017: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 207-230.

Janssen, E., D. J. Wuebbles, K. E. Kunkel, S. C. Olsen, and A. Goodman, 2014: Observational- and model-based trends and projections of extreme precipitation over the contiguous United States. *Earths Future*, **2**, 99-113.

Kunkel, K. E., and Coauthors, 2013: Monitoring and Understanding Trends in Extreme Storms: State of Knowledge. *Bulletin of the American Meteorological Society*, **94**, 499-514.

Mishra, A. K., and V. P. Singh, 2010: Changes in extreme precipitation in Texas. *Journal of Geophysical Research-Atmospheres*, **115**.

^{xxv} Easterling, D. R., and Coauthors, 2017: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 207-230.

^{xxvi} One year block maxima of 1-day precipitation totals for composite stations in each county were fit to a GEV distribution with linearly time-dependent location and scale parameters using the climetRemes Python library. The trend in frequency of extreme rainfall was calculated using the time-dependent PDF, with CMIP5 historical and RCP 4.5 projected global mean surface temperature as a covariate. Citation for software package:

Paciorek, C. J., D. A. Stone, and M. F. Wehner, 2018: Quantifying statistical uncertainty in the attribution of human influence on severe weather. *Weather and Climate Extremes*, **20**, 69-80.

^{xxvii} Trenberth, K. E., A. Dai, R. M. Rasmussen, and D. B. Parsons, 2003: The Changing Character of Precipitation. *Bulletin of the American Meteorological Society*, **84**, 1205-1218.

^{xxviii} Pfahl, S., P. A. O’Gorman, and E. M. Fischer, 2017: Understanding the regional pattern of projected future changes in extreme precipitation. *Nature Climate Change*, **7**, 423.

^{xxix} Easterling, D. R., and Coauthors, 2017: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 207-230.

^{xxx} Based on the 2035 global temperature projection by Kirtman et al. (0.4 °C increase by 2036 compared to 2010-2018) and the ratio of Gulf of Mexico to global temperature increases by Collins et al. (75%-100%).

Kirtman, B., and Coauthors, 2013: Near-term Climate Change: Projections and Predictability. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker, and Coauthors, Eds., Cambridge University Press.

Capotondi, A., and Coauthors, 2015: Understanding ENSO Diversity. *Bulletin of the American Meteorological Society*, **96**, 921-938, Collins, M., and Coauthors, 2013: Long-term Climate Change: Projections, Commitments and Irreversibility. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker, and Coauthors, Eds., Cambridge University Press.

^{xxxii} Based on the county precipitation station analysis, the odds of extreme (once per decade or so) one-day precipitation decrease by a factor of two for every 20% increase in precipitation amount.

^{xxxiii} Based on the updated NOAA Atlas 14 values for Texas, which are used for flood risk calculations.

Perica, S., S. Pavlovic, M. St. Laurent, C. Trypaluk, D. Unruh, and O. Wilhite, 2018: NOAA Atlas 14: Precipitation-Frequency Atlas of the United States, Volume 11 Version 2.0: Texas, 40 pp.

^{xxxiii} Online data archive: <ftp://ftp.ncdc.noaa.gov/pub/data/cirs/climdiv/>

Vose, R. S., and Coauthors, 2014: Improved historical temperature and precipitation time series for U.S. climate divisions. *Journal of Applied Meteorology and Climatology*, **53**, 1232-1251.

^{xxxiv} Data and analysis archive label: climdiv-sp09st-v1.0.0-20190204 and SP09

Online data archive: <ftp://ftp.ncdc.noaa.gov/pub/data/cirs/climdiv/>

Overall yearly drought severity is computed by summing the negative statewide 9-month SPI values for a given year and dividing by 12 to obtain an average annual drought intensity.

^{xxxv} The historic simulations and projections are based on the CMIP5 multi-model ensemble output, as with Footnote iv for temperature, with the following differences: Model simulations using the RCP8.5 scenario are extracted, to improve signal-to-noise ratio. Precipitation is summed for each water year, October-September, rather than each calendar year, in order to better reflect the potential for dryness during the growing season and so as to not divide cool-season precipitation variability associated with El Niño across two separate years. The water year precipitation is converted to anomalies by subtracting the centered 29-year mean for each simulation, with the 1950-1978 mean used for calculating anomalies in 1950-1964. Interannual precipitation variance is then calculated using the same 29-year intervals, and variance time series are created from the overlapping 29-year periods.

^{xxxvi} Pendergrass, A. G., R. Knutti, F. Lehner, C. Deser, and B. M. Sanderson, 2017: Precipitation variability increases in a warmer climate. *Sci Rep*, **7**, 17966.

^{xxxvii} Nielsen-Gammon, J. W., and Coauthors, 2020: Unprecedented Drought Challenges for Texas Water Resources in a Changing Climate: What Do Researchers and Stakeholders Need to Know? *Earth's Future*, **8**, e2020EF001552.

^{xxxviii} Mukherjee, S., A. Mishra, and K. E. Trenberth, 2018: Climate Change and Drought: a Perspective on Drought Indices. *Current Climate Change Reports*, **4**, 145-163.

^{xxxix} Data and analysis archive label: climdiv-pdsist.v1.0.-0-20190204

Online data archive: <ftp://ftp.ncdc.noaa.gov/pub/data/cirs/climdiv/>

Vose, R. S., and Coauthors, 2014: Improved historical temperature and precipitation time series for U.S. climate divisions. *Journal of Applied Meteorology and Climatology*, **53**, 1232-1251.

^{xl} Milly, P. C. D., and K. A. Dunne, 2016: Potential evapotranspiration and continental drying. *Nature Climate Change*, **6**, 946-+.

^{xli} Swann, A. L. S., F. M. Hoffman, C. D. Koven, and J. T. Randerson, 2016: Plant responses to increasing CO2 reduce estimates of climate impacts on drought severity. *Proceedings of the National Academy of Sciences of the United States of America*, **113**, 10019-10024.

Swann, A. L. S., 2018: Plants and Drought in a Changing Climate. *Current Climate Change Reports*, **4**, 192-201. Scheff, J. Ibid.: Drought Indices, Drought Impacts, CO2, and Warming: a Historical and Geologic Perspective, 202-209.

^{xlii} Dai, A., T. Zhao, and J. Chen. Ibid.: Climate Change and Drought: a Precipitation and Evaporation Perspective, 301-312.

Berg, A., and Coauthors, 2016: Land-atmosphere feedbacks amplify aridity increase over land under global warming. *Nature Climate Change*, **6**, 869-+.

^{xliii} Swann, A. L. S., 2018: Plants and Drought in a Changing Climate. *Current Climate Change Reports*, **4**, 192-201.

^{xliv} Cook, B. I., T. R. Ault, and J. E. Smerdon, 2015: Unprecedented 21st century drought risk in the American Southwest and Central Plains. *Science Advances*.

- ^{xlv} Seager, R., A. Hooks, A. P. Williams, B. Cook, J. Nakamura, and N. Henderson, 2015: Climatology, Variability, and Trends in the U.S. Vapor Pressure Deficit, an Important Fire-Related Meteorological Quantity. *Journal of Applied Meteorology and Climatology*, **54**, 1121-1141.
- ^{xlvi} Basis for estimate: Given a statewide average 1 hPa increase in vapor pressure deficit from 1960-2013 and a climatological mean warm season vapor pressure deficit of 14 hPa (both from Seager et al. 2015), the fractional trend is about -1/14 in 50 years, or -1/7 in 100 years. Over the 26 or so years from 2000-2018 to 2036, a continuation of the trend would yield a change of about -1/28, or -4%.
- ^{xlvii} Berg, A., J. Sheffield, and P. C. D. Milly, 2017: Divergent surface and total soil moisture projections under global warming. *Geophysical Research Letters*, **44**, 236-244.
- ^{xlviii} Pendergrass, A. G., R. Knutti, F. Lehner, C. Deser, and B. M. Sanderson, 2017: Precipitation variability increases in a warmer climate. *Sci Rep*, **7**, 17966.
- ^{xlix} Bukovsky, M. S., R. R. McCrary, A. Seth, and L. O. Mearns, 2017: A Mechanistically Credible, Poleward Shift in Warm-Season Precipitation Projected for the U.S. Southern Great Plains? *Journal of Climate*, **30**, 8275-8298.
- ^l Qing, Y., S. Wang, B. Zhang, and Y. Wang, 2020: Ultra-high resolution regional climate projections for assessing changes in hydrological extremes and underlying uncertainties. *Climate Dynamics*, **55**, 2031-2051.
The paper appears to have calculated SPI and related indices without taking into account the annual cycle of precipitation, temperature, and other parameters.
- ^{li} Cleaveland, M. K., T. H. Votteler, D. K. Stahle, R. C. Casteel, and J. L. Banner, 2011: Extended Chronology of Drought in South Central, Southeastern and West Texas. *Texas Water Journal*, **2**, 54-96.
- ^{lii} Ashley, S. T., and W. S. Ashley, 2008: Flood Fatalities in the United States. *Journal of Applied Meteorology and Climatology*, **47**, 805-818.
- ^{liii} Berghuijs, W. R., R. A. Woods, C. J. Hutton, and M. Sivapalan, 2016: Dominant flood generating mechanisms across the United States. *Geophysical Research Letters*, **43**, 4382-4390.
- ^{liv} Villarini, G., and J. A. Smith, 2013: FLOODING IN TEXAS: EXAMINATION OF TEMPORAL CHANGES AND IMPACTS OF TROPICAL CYCLONES. *Journal of the American Water Resources Association*, **49**, 825-837.
- ^{lv} Mishra, A. K., V. P. Singh, and M. Özger, 2011: Seasonal streamflow extremes in Texas river basins: Uncertainty, trends, and teleconnections. *Journal of Geophysical Research*, **116**.
- ^{lvi} Qing, Y., S. Wang, B. Zhang, and Y. Wang, 2020: Ultra-high resolution regional climate projections for assessing changes in hydrological extremes and underlying uncertainties. *Climate Dynamics*, **55**, 2031-2051.
- ^{lvii} Li, X., and Coauthors, 2020: Impacts of urbanization, antecedent rainfall event, and cyclone tracks on extreme floods at Houston reservoirs during Hurricane Harvey. *Environmental Research Letters*, **15**, 124012.
- ^{lviii} Berg, M. D., 2018: Peak flow trends highlight emerging urban flooding hotspots in Texas. *Texas Water Journal*, **9**, 18-29.
- ^{lix} Zhang, W., G. Villarini, G. A. Vecchi, and J. A. Smith, 2018: Urbanization exacerbated the rainfall and flooding caused by hurricane Harvey in Houston. *Nature*, **563**, 384-388.
Among our concerns with this study are the use of generic building heights rather than actual lidar-measured heights to represent Houston urbanization and a method of aggregating streamflow trends that gives undue influence to outliers.
- ^{lx} Berg, M. D., 2018: Peak flow trends highlight emerging urban flooding hotspots in Texas. *Texas Water Journal*, **9**, 18-29, Villarini, G., and J. A. Smith, 2013: FLOODING IN TEXAS: EXAMINATION OF TEMPORAL CHANGES AND IMPACTS OF TROPICAL CYCLONES. *Journal of the American Water Resources Association*, **49**, 825-837.
- ^{lxi} Data accessed from U.S. Census Bureau, <https://www.census.gov/geo/reference/ua/urban-rural-2010.html>, 18 February 2019.
- ^{lxii} Zhao, G., H. L. Gao, and L. Cuo, 2016: Effects of Urbanization and Climate Change on Peak Flows over the San Antonio River Basin, Texas. *Journal of Hydrometeorology*, **17**, 2371-2389.
- ^{lxiii} The county-scale historic snowfall analysis is generated from county-specific composite station data, as with the county-scale temperature and precipitation analyses discussed earlier in the report, except that the number of allowed missing data points per year is increased to 27 to allow for the lower reliability of reporting of null snowfall amounts. Also, for snowfall a year is defined as beginning on July 1 and ending on June 30.
- ^{lxiv} Lanza, M., 2017: Space City Rewind: Houston's Great Snow of 1895. *Space City Weather*. [Accessed March 16, 2019]
- ^{lxv} Bomar, G. W., 2017: *Weather in Texas: The Essential Handbook*. Third Edition ed. University of Texas Press, 290 pp, TSHA, 2018: *Texas Almanac 2018-2019*. Texas State Historical Association, 752 pp.

- ^{lxvi} Kluver, D., and D. Leathers, 2015: Winter snowfall prediction in the United States using multiple discriminant analysis. *International Journal of Climatology*, **35**, 2003-2018.
- ^{lxvii} O’Gorman, P. A., 2014: Contrasting responses of mean and extreme snowfall to climate change. *Nature*, **512**, 416-418.
- ^{lxviii} Changnon, S. A., and T. R. Karl, 2003: Temporal and Spatial Variations of Freezing Rain in the Contiguous United States: 1948–2000. *Journal of Applied Meteorology*, **42**, 1302-1315.
- ^{lxix} Data source: Storm Prediction Center, National Oceanic and Atmospheric Administration, <https://www.spc.noaa.gov/gis/svrgis/>
- ^{lxx} For example, see:
Anderson, C. J., C. K. Wikle, Q. Zhou, and J. A. Royle, 2007: Population Influences on Tornado Reports in the United States. *Weather and Forecasting*, **22**, 571-579.
Elsner, J. B., L. E. Michaels, K. N. Scheitlin, and I. J. Elsner, 2013: The Decreasing Population Bias in Tornado Reports across the Central Plains. *Weather, Climate, and Society*, **5**, 221-232.
- ^{lxxi} Taszarek, M., J. T. Allen, H. E. Brooks, N. Pilguy, and B. Czernecki, 2021: Differing Trends in United States and European Severe Thunderstorm Environments in a Warming Climate. *Bulletin of the American Meteorological Society*, **102**, E296-E322.
- Koch, E., J. Koh, A. C. Davison, C. Lepore, and M. K. Tippett, 2021: Trends in the Extremes of Environments Associated with Severe U.S. Thunderstorms. *Journal of Climate*, **34**, 1259-1272.
- Tang, B. H., V. A. Gensini, and C. R. Homeyer, 2019: Trends in United States large hail environments and observations. *npj Climate and Atmospheric Science*, **2**, 45.
- ^{lxxii} Van Klooster, S. L., and P. J. Roebber, 2009: Surface-Based Convective Potential in the Contiguous United States in a Business-as-Usual Future Climate. *Journal of Climate*, **22**, 3317-3330.
- ^{lxxiii} Brooks, H. E., 2013: Severe thunderstorms and climate change. *Atmospheric Research*, **123**, 129-138,
Difffenbaugh, N. S., M. Scherer, and R. J. Trapp, 2013: Robust increases in severe thunderstorm environments in response to greenhouse forcing. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, 16361-16366.
- ^{lxxiv} Smith, B. T., T. E. Castellanos, A. C. Wintners, C. M. Mead, A. R. Dean, and R. L. Thompson, 2013: Measured Severe Convective Wind Climatology and Associated Convective Modes of Thunderstorms in the Contiguous United States, 2003–09. *Weather and Forecasting*, **28**, 229-236.
- ^{lxxv} Tippett, M. K., C. Lepore, and J. E. Cohen, 2016: More tornadoes in the most extreme U.S. tornado outbreaks. *Science*, **354**, 1419-1423.
- ^{lxxvi} Brimelow, J. C., W. R. Burrows, and J. M. Hanesiak, 2017: The changing hail threat over North America in response to anthropogenic climate change. *Nature Climate Change*, **7**, 516.
- Tang, B. H., V. A. Gensini, and C. R. Homeyer, 2019: Trends in United States large hail environments and observations. *npj Climate and Atmospheric Science*, **2**, 45.
- ^{lxxvii} Tide gauge data is from the National Oceanic and Atmospheric Administration, <https://tidesandcurrents.noaa.gov/sltrends/>
- ^{lxxviii} Letetrel, C., M. Karpytchev, M. N. Bouin, M. Marcos, A. Santamaría-Gómez, and G. Wöppelmann, 2015: Estimation of vertical land movement rates along the coasts of the Gulf of Mexico over the past decades. *Continental Shelf Research*, **111**, 42-51.
- ^{lxxix} Davis, R. A. J., 2011: *Sea-Level Change in the Gulf of Mexico*. Texas A&M University Press, 171 pp.
- ^{lxxx} Paine, J. G., T. L. Caudle, and J. R. Andrews, 2017: Shoreline and Sand Storage Dynamics from Annual Airborne LIDAR Surveys, Texas Gulf Coast. *Journal of Coastal Research*, **33**, 487-506.
- Paine, J. G., and T. L. Caudle, 2020: Shoreline Movement along the Texas Gulf Coast, 1930s to 2019, 64 pp.
- ^{lxxxii} Al Mukaimi, M. E., T. M. Dellapenna, and J. R. Williams, 2018: Enhanced land subsidence in Galveston Bay, Texas: Interaction between sediment accumulation rates and relative sea level rise. *Estuarine, Coastal and Shelf Science*, **207**, 183-193.
- ^{lxxxiii} Frederikse, T., S. Jevrejeva, R. E. M. Riva, and S. Dangendorf, 2018: A Consistent Sea-Level Reconstruction and Its Budget on Basin and Global Scales over 1958-2014. *Journal of Climate*, **31**, 1267-1280.
- Nerem, R. S., B. D. Beckley, J. T. Fasullo, B. D. Hamlington, D. Masters, and G. T. Mitchum, 2018: Climate-change-driven accelerated sea-level rise detected in the altimeter era. *Proceedings of the National Academy of Sciences of the United States of America*, **115**, 2022-2025.
- Dangendorf, S., C. Hay, F. M. Calafat, M. Marcos, C. G. Piecuch, K. Berk, and J. Jensen, 2019: Persistent acceleration in global sea-level rise since the 1960s. *Nature Climate Change*, **9**, 705-710.

- ^{lxxxiii} Kossin, J. P., T. Hall, T. R. Knutson, K. E. Kunkel, R. J. Trapp, D. E. Waliser, and M. E. Wehner, 2017: Extreme Storms. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock, Eds., U.S. Global Change Research Program, 257-276, Needham, H. F., B. D. Keim, D. Sathiaraj, and M. Shafer, 2012: Storm Surge Return Periods for the United States Gulf Coast. *Advances in Hurricane Engineering*, 715-740.
- ^{lxxxiv} Knutson, T., and Coauthors, 2019: Tropical Cyclones and Climate Change Assessment: Part I: Detection and Attribution. *Bulletin of the American Meteorological Society*, **100**, 1987-2007.
- , 2020: Tropical Cyclones and Climate Change Assessment: Part II: Projected Response to Anthropogenic Warming. *Bulletin of the American Meteorological Society*, **101**, E303-E322.
- ^{lxxxv} Hall, T. M., and J. P. Kossin, 2019: Hurricane stalling along the North American coast and implications for rainfall. *npj Climate and Atmospheric Science*, **2**, 17.
- Hassanzadeh, P., C.-Y. Lee, E. Nabizadeh, S. J. Camargo, D. Ma, and L. Y. Yeung, 2020: Effects of climate change on the movement of future landfalling Texas tropical cyclones. *Nature Communications*, **11**, 3319.
- ^{lxxxvi} Camargo, S. J., and A. A. Wing, 2016: Tropical cyclones in climate models. *Wiley Interdisciplinary Reviews: Climate Change*, **7**, 211-237.
- ^{lxxxvii} Pryor, S. C., and J. Ledolter, 2010: Addendum to “Wind speed trends over the contiguous United States”. *Journal of Geophysical Research: Atmospheres*, **115**.
- ^{lxxxviii} Liu, Y., S. L. Goodrick, and J. A. Stanturf, 2013: Future U.S. wildfire potential trends projected using a dynamically downscaled climate change scenario. *Forest Ecology and Management*, **294**, 120-135.
- ^{lxxxix} Huang, Y., S. Wu, and J. O. Kaplan, 2015: Sensitivity of global wildfire occurrences to various factors in the context of global change. *Atmospheric Environment*, **121**, 86-92.
- ^{xc} Liu, Y., S. L. Goodrick, and J. A. Stanturf, 2013: Future U.S. wildfire potential trends projected using a dynamically downscaled climate change scenario. *Forest Ecology and Management*, **294**, 120-135.
- ^{xci} Guyette, R. P., F. R. Thompson, J. Whittier, M. C. Stambaugh, and D. C. Dey, 2014: Future Fire Probability Modeling with Climate Change Data and Physical Chemistry. *Forest Science*, **60**, 862-870, Stambaugh, M. C., R. P. Guyette, E. D. Stroh, M. A. Struckhoff, and J. B. Whittier, 2018: Future southcentral US wildfire probability due to climate change. *Climatic Change*, **147**, 617-631.







TEXAS A&M UNIVERSITY

College of Geosciences

